

Brown Marmorated Stink Bug:

Yet another new kid on the block...



Christelle Guédot


UW Department of Entomology

guedot@wisc.edu

(608) 262-0899

Appearance

Hemiptera: Pentatomidae: *Halyomorpha halys*



- Mottled brown to grey
- $\frac{1}{2}$ - $\frac{3}{4}$ " long
- Legs are brown and may have faint white bands

BMSB look-alikes



Acrosternum sp.




Euschistus sp.



Boxelder Bug.



Western Conifer Seed Bug



BMSB



Brochymena

Confirmation

- If you find or suspect BMSB, please send specimen in alcohol to:
Insect Diagnostic Lab
1630 Linden Dr.
University of Wisconsin
Madison, WI 53706
- Rutgers University has an active monitoring program for the spread of BMSB
- If you find BMSB, please fill-out and submit a secure, on-line form at: <https://njaes.rutgers.edu/stinkbug/report.asp>

Life cycle

Eggs



- ~0.06"
- Clusters of 20-30
- Underside of leaves

1st instars



- Bright orange/red
- Black head and thorax
- emerge in 4-5 days

2nd instars



- Very dark
- Tick-like

Life cycle

3rd to 5th instars



Images from: bugguide.net; Kenneth Frank



Each instar lasts ~1 week

- Reddish to off-white abdomen
- Dark markings center of back and along margins of abdomen
- White bands on legs and antennae

Biology

- Adults emerge late March-June
- Adults sexually mature in ~2 weeks
- Each female can lay about 250 eggs
- Nymphs tend to be solitary feeders, but often congregate on leaves, bark, or fruit
- New generation of adults begin to appear in mid to late summer



Photo credit Mike Raupp

Biology

- Probably would be only 1 generation in WI (1 in NJ, 2 in WV)
- Feeding increases at night (0-4AM)



Overwintering and pest status

- Adults overwinter in protected locations - natural rocky outcroppings, houses and structures, in bark of dead standing trees
- Adults sheltering in homes active on warm days in late winter (active in March this year in Madison!)



Overwintering

- Adults overwinter in large (>7.5" dbh) dry dead standing trees with loose and peeling bark
- Hide beneath bark or inside decomposed woody tissue
- Dogs used for detection!
- Favored hosts: oak, locust, tree of heaven, elm
- Stink bugs found in leaf litter, but no BMSB
- Edge effect and early season hosts surrounding crops (maple, catalpa, redbud, walnut, mulberry, cherry, black locust...)


Introduction

Native from Asia (China, Korea, Japan)

Probably came to US on shipping boat in 1996 (Allentown, PA)




Potential geographic distribution



Dark color represents high suitability


Potential geographic distribution



Dark color represents high suitability

Actual distribution

As of December 2015, BMSB has been detected in 42 states



Michigan in 2010
2015 moved to Ag
problem


Distribution in Wisconsin

2010: First detection (car)

2011: First report in Dane and Jefferson counties

2012: Breeding populations reported in several counties

2015: No nuisance or agricultural problem reported, but numbers increasing in urban areas



Host plants

Significant agricultural pest in some areas of the eastern US

- Tree fruits: apple, peach, cherry, pear,...
- Vegetables: peppers, tomatoes, asparagus...
- Fruits: berries, grapes, currant, melons,...
- Field crops: soybeans, corn,...
- Trees: maple, crabapple, oak, black cherry

(feeds through the bark)



Photo credit Mike Raupp

Management options

- Population monitoring and trapping
- Chemical control
- Biological control


Population monitoring

- Scouting: edge effect, higher abundance in border rows, adjacent to woods especially
- Research on border row sprays in NJ
- Season-long monitoring now possible with pheromone traps
- Light traps also tested


BMSB chemical attractants

BMSB aggregation pheromone and *Plautia stali* aggregation pheromone

Main component of BMSB aggregation pheromone
(3*S*,6*S*,7*R*,10*S*)-10,11-epoxy-1-bisabolen-3-ol




Minor component of BMSB aggregation pheromone
(3*R*,6*S*,7*R*,10*S*)-10,11-epoxy-1-bisabolen-3-ol



+

Methyl (*E,E,Z*)-2,4,6-decatrienoate (MDT) acts as a synergist for BMSB pheromone




=

Synergism

Slide courtesy of Dr. Tracy Leskey, USDA ARS
Greg Krawczyk, PSU FREC 2014


BMSB chemical attractants

Synergism of pheromone and MDT works all season and for adults and nymphs



BMSB chemical attractants

Proportion of trap catches throughout season



BMSB traps



Dead-Inn pyramid trap
Ag Bio
\$35/trap



Experimental trap
USDA-ARS



RESCUE® Stink Bug Trap
Great Lakes IPM
\$20/trap with attractant
Light attachment \$17.30



DEAD-INN Stink Bug Trap and Box
Creative Commons from fauxto_digit
\$24.95 from EcoGreen Warehouse



Lure from Ag Bio
\$6 per lure

BMSB light traps



Experimental trap
USDA-ARS




Night View



BMSB light traps

Tests of different light waves and frequencies for BMSB specificity



Chemical control

Class	Trade name	Active ingredient	PHI (days)	Probable efficacy
Neonicotinoid	Actara	Thiamethoxam	35	+++
	Belay	Clothianidin	7	+++++
Pyrethroid	Baythroid	β -cyfluthrin	7	+++
	Danitol	Fenpropathrin	14	+++
Carbamate	Lannate	Methomyl	14	++
Organochlorin	Endosulfan	Endosulfan	21	+++

Follow the label, it is the law!!

Biological control

Native parasitoids and predators:

- *Telenomus podisi* (Hymenoptera: Scelionidae) native to U.S.
- *Trissolcus japonicus*: egg parasitoid, primary BC agent responsible for management of BMSB in northern China: 50% mortality rate for BMSB populations. Not known in U.S.
- Predation of eggs in U.S. by Carabidae and Harpalus (ground beetles), Salticids (spiders), earwigs, Tettigoniidae (bush crickets).



Telenomus podisi



Trissolcus japonicus

Biological control

<https://www.youtube.com/watch?v=rbdXiiM538I>



A photograph of a tree-lined path in a park, with the word 'QUESTIONS???' overlaid in large white text.

QUESTIONS???

Christelle Guédot

Email: guedot@wisc.edu

Phone: (608) 262-0899